Did your doctor tell you that you have developed an infection related to your surgery?

If YES, please call the Infection Prevention Control office at LWDH

(807) 468-9861 extension 2362

Thank you for helping us ensure quality care.

Lake of the Woods District Hospital

21 Sylvan Street Kenora, Ontario P9N 3W7 Telephone (807)468-9861

DISCHARGE INSTRUCTIONS

FOR

LAPAROTOMY

Discharge Instructions: Laparotomy 405026 09/06/2019

- Consume a diet high in protein to promote healing. (if able to) If you feel uncomfortable, stay on clear fluids until the feeling passes. If nauseated, avoid milk products until settled.
- 2. Increase intake of fluid and fiber to avoid constipation. Laxatives may be used for constipation but please discuss this with your surgeon first.
- 3. **Do not lift anything heavier than 10 pounds** (6 kg) for 6 weeks or as otherwise directed by your doctor.
- 4. Avoid having a bath until steri-strips, staples and drainage tubes are removed. **You may shower.** When showering, avoid saturating the incision with water while steri-strips are in place. Pat the incision dry. Avoid using soaps, powders and lotions.
- 5. You may leave your incision exposed to air or cover with a light dressing for comfort.
- 6. Resume medications as before your procedure unless otherwise advised.
- 7. Resume normal activity as tolerated and as directed.
- 8. Take your prescribed pain medication as directed for discomfort.
- 9. Your physician will advise you when you may return to work.
- 10. If you have steri-strips along your incision, you may remove them in 1 week.

11. If you have been given anesthesia or sedation;

- Do not drive or operate machinery for 24 hours.
- Do not make important decisions or sign any important documents in the next 24 hours. Do not plan any activities that require complex thought or coordination for 24 hours.
- Do not consume alcohol, tranquilizers or sleeping medications for the next 24 hours.
- Sedative and narcotic medicines given during a surgical procedure may linger for hours. You may feel dizzy for several hours after your procedure.
- You must have an adult with you tonight at home.
- 12. **Go to an Emergency Department** if any of the following occurs:
 - Excessive pain not relieved by medications.
 - Redness, swelling or drainage of pus from incision
 - Elevated temperature (over 38°C or 100°F)
 - Persistent nausea or vomiting not relieved by over the counter medications.
 - Bladder difficulties
 - Unusual bleeding
 - Unusual cough or shortness of breath

13. Make an a	ppointment to see Dr
in	Phone number
	If you have any questions, do not hesitate to call your doctor.
Special Instruc	etions:
	

Discharge Instructions: Laparotomy 405026 09/06/2019